

Environmental Hazards

Policy

“The anesthetic gases and vapors that leak into the surrounding room during medical procedures are considered waste anesthetic gases. Health care professionals who work in hospitals, operating rooms, dental offices and veterinary clinics, are potentially exposed to waste anesthetic gases and are at risk of occupational illness. The waste anesthetic gases and vapors of concern are nitrous oxide and halogenated agents (vapors) such as halothane, enflurane, isoflurane, and desflurane. Some potential effects of exposure to waste anesthetic gases are nausea, dizziness, headaches, fatigue, and irritability, as well as sterility, miscarriages, birth defects, cancer, and liver and kidney disease, among operating room staff or their spouses (in the case of miscarriages and birth defects). Employers and employees should be aware of the potential effects and be advised to take appropriate precautions.”

[OSHA Waste Anesthetic Gases](#)

[OSHA Reproductive Hazards](#)

Selected inhalation anesthetic agents are thought to be hepatotoxic and, as a result, an anesthetist may on occasion develop a sensitivity to agents which is reflected in abnormal liver function studies. In addition, studies performed in the past have suggested an association between sustained exposure to an anesthetic environment and an increased incidence in abortions, birth defects, and certain types of malignancies for both male and female personnel. While no cause-and-effect relationship has been established, consideration should be given to these findings in choosing anesthesia as a specialty.

Most hospitals have installed anesthetic gas exhaust systems to minimize risk to operating room personnel. All clinical affiliates have scavenging systems for waste gases. It has not been established whether any risks to personnel are eliminated by these exhaust systems.

Anesthesia providers are frequently exposed to blood products, body secretions, and contaminated syringes and needles. All nurse anesthesia residents must strictly adhere to universal precautions when involved in patient care involving potential contact with mucous membranes, secretions or open wounds. Gloves, protective eye wear and masks are available at each anesthetizing site and must be worn. Proper regard for and performance of aseptic techniques is mandatory to protect both patients and anesthesia providers. All residents must have current vaccinations including those for Hepatitis B or have appropriate titer levels. If a resident experiences a needle stick with a contaminated needle they should:

1. Notify the Clinical Coordinator
2. Seek medical attention per the clinical site's policy
3. Complete required hospital forms
4. Notify the Program Director
5. Related medical expenses accrued by the resident, not covered by the resident's health insurance, are the responsibility of the resident.

Anesthesia personnel are also frequently exposed to x-rays during operative procedures. Lead aprons and thyroid shields are available at each anesthetizing site and must be worn during fluoroscopy and x-ray procedures. Radiology dosimeters to monitor the level of x-ray exposure may be properly worn.

Responsibility for accepting risks associated with this specialty rests with the individual who chooses to work within this environment, rather than with the institutions which take reasonable precautions to minimize potential hazards.

[CDC Other Healthcare Worker Hazards](#)

Written: 11-09; Reviewed and Revised (R&R) 8-12; R&R June 2014; Reviewed June 2016; Reviewed July 2017; R&R July 2018; Reviewed 7-19; Revised 8-20; R&R Aug 2021, Reviewed December 2022; R&R March 2024; Reviewed May 2025; R&R April 2026